



Florian Luca

University of the Witwatersrand

Fecha: Miércoles 22 de noviembre 2017

Hora: 14:00-15:00

Lugar: Universidad de Valparaíso,
Facultad de Ciencias, sala 12

Title: Cyclotomic factors of Serre's polynomials

Abstract: Consider the family of polynomials $P_m(X) \in \mathbb{Q}[X]$ given by

$$\prod_{m \geq 1} (1 - q^m)^{-z} = \sum_{m \geq 0} P_m(z) q^m.$$

These polynomials have deep connections with the theory of partition numbers and the Ramanujan τ -function. They appeared for the first time in work of Newman 1955, and were used by Serre in his 1985 work on the lacunarity of the powers of the Dedekind eta function. They can also be given recursively as $P_0(X) = 1$ and

$$P_m(X) = \frac{X}{m} \left(\sum_{k=1}^m \sigma(k) P_{m-k}(X) \right).$$

It is easy to see that $P_m(X)$ has no positive real roots. Further, by the Euler pentagonal formula, it follows that $X + 1 \mid P_m(X)$ for infinitely many m . We ask whether $P_m(X)$ can have other roots of unity except -1 . We prove that this is never the case, namely that if ζ is a root of unity of order $N \geq 3$ and $m \geq 1$, then $P_m(\zeta) \neq 0$. The proof uses basic facts about finite fields and a bit of analytic number theory.

This is joint work with: Bernhard Heim and Markus Neuhauser.

<http://seminarioaritméticaygeometría.wordpress.com>